

## **Social enrichment by separated pair housing of male C57BL/6JRj mice**

Hohlbaum, K<sup>1</sup>, Frahm, S<sup>2</sup>, Rex, A<sup>3</sup>, Palme, R<sup>4</sup>, Thöne-Reineke, C<sup>1</sup>, Ullmann, K<sup>5, 6\*</sup>

<sup>1</sup> Institute of Animal Welfare, Animal Behavior, and Laboratory Animal Science, Department of Veterinary Medicine, Freie Universität Berlin, Berlin, Germany

<sup>2</sup> Department of Pharmacology, Charité – Universitätsmedizin Berlin, Berlin, Germany

<sup>3</sup> Department of experimental Neurology, Charité - Universitätsmedizin Berlin, Berlin, Germany

<sup>4</sup> Unit of Physiology, Pathophysiology and Experimental Endocrinology, Department of Biomedical Sciences, University of Veterinary Medicine, Vienna, Austria

<sup>5</sup> FEM, Charité - Universitätsmedizin Berlin, Berlin, Germany

<sup>6</sup> Charité 3R, Charité - Universitätsmedizin Berlin, Berlin, Germany

\* Corresponding author: [kristina.ullmann@charite.de](mailto:kristina.ullmann@charite.de)

## Abstract

Laboratory male mice are often housed individually due to aggressions or experimental requirements, though social isolation can cause welfare issues. As a strategy to refine housing of male mice, we introduce the separated pair housing system. A perforated transparent wall divides the cage into two compartments and allows olfactory, acoustical, and visual communication between the two mice but prevents fighting and injuries. Long-term effects of separated pair housing on well-being and distress of adult male C57BL/6JRj mice were investigated and compared with both single and group housed mice. Behavioral analysis after eight weeks in three different housing systems revealed no differences in burrowing performance, social interaction, anxiety, and stress hormone concentrations. However, pair-housed mice built more complex nests compared to single-housed mice and the nest position suggested that pair-housed mice preferred the close proximity to their cage mates. Moreover, pair-housed mice showed less locomotor activity compared to group- and single-housed mice. Body weight was higher in group-housed mice. All in all, no unambiguous long-term beneficial effects of pair housing on the well-being were found. However, the findings emphasized that effects of the housing systems on behavioral, physical, and biochemical parameters must be considered in the design of animal experimental studies.

## Introduction

In nature, wild male mice live as loners or in polygamous family groups[1]. To avoid uncontrolled breeding in laboratory animal facilities, it is a common code of practice to house laboratory mice in single-sex groups [2-4]. However, inter-individual aggression can result in stress or, at worst, pain, injuries or death [5-7]. Although there are some approaches how to decrease aggressive behaviors in male mice, e.g. by partial cage division [8], they often have to be separated and housed in isolation when the aggression behavior exceeds an unacceptable level (e.g. fights, wounds, body weight loss) [7]. Besides the prevention of aggression, further justifications for housing mice individually are veterinary needs and experimental requirements [2,5]. In comparison to group-housed mice, individually housed mice show lower variances in physiological parameters like body fat and bone mineral content [9] and adapt faster to stress induced by cage cleaning [10]; nevertheless, single housing can also diminish well-being of mice. Social isolation is associated with depressive states [11-13], lower tolerance to external stressors [14], increased stress-related corticosterone concentrations [15], and varying occurrence of anxiety-related behavior [13,16,17] in mice.

In the scope of refinement, single housing “should be limited to the minimum period necessary, and where possible, visual, auditory, olfactory, and tactile contact with compatible conspecifics should be provided” [2]. If individually housed mice from adult age have the choice between an empty cage or a cage inhabited by another male, they prefer the company of the other mouse [18,19]. Therefore, one strategy to ameliorate the negative effects of single housing on the well-being of mice is to allow the animals social contact with conspecifics. A cage divider, e.g. a grid, separating the cage into two compartments prevents fighting and injuries but makes sensory contact possible. Short-term effects of this housing condition (hereafter referred to as separated pair housing) have already been investigated: heart rate, body temperature, motor activity, body weight and nest building behavior were altered in 8–9-month-old vasectomized Hsd:NMRI mice when being housed separated by a

grid with sensory contact to an unfamiliar male for 10 days [20]. Moreover, normalization of heart rate was delayed after the end of this observation period [20]. Although these effects indicated social stress and, subsequently, reduced animal well-being during the habituation to the new housing condition, long-term effects are unknown. The social stress may deteriorate, but it is also possible that mice get used to the new cage mate living beyond the grid and benefit from its company. To investigate whether this housing condition can foster well-being of male mice, we compared long-term effects of housing adult male C57BL/6JRj mice in pairs, separated by a perforated transparent wall, which allows olfactory, acoustical, and visual communication, to single and group housed mice. We systematically assessed well-being of the mice using physical, biochemical, and behavioral parameters, whereby any additional stress caused by the test battery was minimized by using non-invasive methods. We analyzed burrowing and nesting, trait anxiety-related behavior in the free exploratory paradigm, the ease of handling, and social behavior in a social interaction test. Additionally, body weight and stress hormone (metabolites) concentrations in feces and hair were measured.

## **Material and Methods**

### **Ethics**

The study was performed according to the guidelines of the German Animal Welfare Act and the Directive 2010/63/EU for the protection of animals used for scientific purposes.

Maintenance of mice and all animal experimentation were approved by the Berlin State Authority (“Landesamt für Gesundheit und Soziales”, permit number: G 0251/18) and registered in Animal Study Registry (DOI: 10.17590/asr.0000101). Sample size calculation (primary outcome measure: effect of the housing systems on hair corticosterone) was performed using package “pwr” in R (power of 80 %, standardized effect size of 0.5).

## **Animals**

A total number of 60 adult male C57BL/6JRj mice obtained from Janvier Labs (Saint-Berthevin Cedex, France) at 4 weeks of age and were allowed to habituate to our animal facility for three weeks. This strain was chosen since C57BL/6JRj mice are the most commonly used laboratory mice. The mice were assigned to three study groups by simple randomization: single housing (n = 16), group housing (n = 16), and pair housing (n = 16). Twelve mice served as target animals in the social interaction test.

## **Housing conditions**

Mice were housed in individually ventilated cages (IVCs). The cages contained wooden bedding material (Safe<sup>R</sup> Select, Safe, Augy, France) and nestlets (Ancare, UK agents, Lillico, United Kingdom) and a red, triangular plastic house (length: 12,5 cm, width: 11 cm, height: 6 cm; Tecniplast, Italy) or a plastic tunnel (length: 10 cm, diameter: 4,5 cm, self-made). The animals were maintained under standard conditions (room temperature:  $22 \pm 2$  °C; relative humidity:  $55 \pm 10$  %) on a light:dark cycle of 12:12 hours of artificial light (lights on from 6:00 a.m. to 6:00 p.m.). The mice were fed pelleted mouse diet ad libitum (V1534-000, Ssniff, Soest, Germany) and had free access to tap water.

During habituation mice were kept in groups of 3–4 siblings per cage in Polysulfone type II long cages (32 cm × 17 cm × 13 cm). Mice assigned to the study group “group housing” stayed in this group (i.e. their group constellation was not changed after the habituation period). Single-housed mice were transferred to Polysulfone type I long cages (32 cm × 13 cm × 13 cm) and pair-housed mice to Green Line IVC Sealsafe PLUS Rat GR 900, which were separated in two compartments (size of one compartment: 31 cm × 17 cm × 15 cm) by a perforated transparent wall (cage divider) allowing olfactory, acoustical, and visual communication (Figure 1). Mice sharing a pair-housing cage had not known each other before.



**Figure 1. Pair housing system** from the side view (a) and top view (b).


### **Testing schedule**

Figure 2 gives an overview of the testing schedule. After mice had habituated to the animal facility for approximately three weeks, they were transferred to the respective housing system. During the first day in the (new) housing system, cages did not contain houses or tunnels because nest complexity was scored on day 1 (7.00 and 7.30 A.M.). Subsequent to nest scoring, mice were moved to the testing room and were allowed to habituate for approximately an hour (~7.30–8.30 A.M.). Then they were transferred to the testing cage and acclimatized for 30 min (~8.30–9 A.M.) to the new environment, before the test for burrowing was carried out (~9.00–11.00 A.M.). After burrows were removed from the testing cages, the free exploratory paradigm for trait anxiety-related behavior was carried out. All fecal pellets mice had excreted during the period they spent in the testing cage were collected. Furthermore, hair samples were taken. Finally, the mice were transferred to their home cages and houses or tunnels were added.

In week 7, the social interaction test was performed and a week later, in week 8, the ease of handling, nesting, burrowing, and trait anxiety-related behavior were tested and fecal as well as hair samples were taken. After week 8, single-housed mice were re-socialized and transferred to the pair-housing system (hereafter also referred to as single/pair housing). In

week 11, all mice were tested again in the social interaction test and, in week 12, the other parameters, except from the ease of handling, were investigated for a third time.

Body weight was determined once a week when cages were changed. All efforts were made to blind experimenters wherever possible. However, unfortunately, the study design did not allow us to blind experimenters for nesting, and the tests for ease of handling because these investigations were carried out in the home cage of the animals.



**Figure 2. Testing schedule.** Behavioral, biochemical, and physical parameters were measured to investigate the well-being of mice kept in single, group, and pair housing.

## Burrowing

The burrowing performance [21,22] was tested by using a blue opaque plastic water bottle as burrow (7 cm x 7 cm x 11.5 cm, 3 cm diameter of bottle neck) which was filled with approximately 140 ± 2 g food pellets normally supplied as diet [23,24]. After a 30-min habituation period to the testing cage (Polysulfone type III cages, 42 x 26 x 15 cm, approximately 0.5–1 cm bedding material, water bottle, feeder filled with a few pellets), the burrow was placed in a corner, parallel to the left long wall of the cage and the mice had been allowed to burrow for two hours. At the end of this test, the weight of food pellets removed from the burrow was determined.

### **Nesting**

When cages were routinely changed, a square cotton nestlet with a weight of 2.4–2.6 g (Ancare, Bellmore, NY, USA) was placed in the front left corner. In the morning of the following day (i.e. approximately 1–1.5 hours after lights turned on), nest complexity was scored on a 6-point-scale using the protocol by Hess et al. [25,26]. Moreover, the position of the nest was noted in pair-housed mice to evaluate whether they prefer distance or proximity to the other mouse.

### **Ease of Handling**

The ease of handling was tested during the weekly cage change in week 8. The voluntary approach and interaction of the mice with the experimenter's hand can provide valuable information on the "anxiety-related behavior in anticipation of handling" [27]. The cage was transferred to a table and the lid, the grid, as well as the house were removed. The left gloved hand (with the palm facing downwards) was placed on the opposite cage site to the nest location. Pair-housed mice were tested simultaneously (i.e. the left hand was used for the left compartment and the right hand for the right compartment). The mice were monitored for 60 seconds and the latency to first voluntary interaction with the experimenter's hand (i.e.



experimenter perceives contact of whiskers, nose, and/or paws with hand) was noted.

Moreover, within the 60-second testing period, their interaction with the hand was scored on a rating scale for voluntary interaction:

- Score 0 = The mouse moved away from the hand and settled down at the largest possible distance. It made no attempts to approach the experimenter's hand.
- Score 1 = The mouse carefully approached but did not touch the experimenter's hand. Protected stretches towards the hand could be observed.
- Score 2 = The mouse explored the experimenter's hand by direct contact of whiskers and/or nose.
- Score 3 = The mouse explored the experimenter's hand by direct contact of paws.
- Score 4 = The mouse explored the experimenter's hand by climbing on it.

After monitoring the interaction of the mouse with the hand, the animal was picked up by the tail and gently transferred to the back of the hand. The process of catching and picking up the mouse was scored on a 5-point scale (rating scale for capture, modified from Wahlsten et al. [28]):

- Score 0 = The experimenter caught the mouse at the first try. The mouse showed minimal resistance to capture.
- Score 1 = The mouse escaped from the first try of the experimenter to catch it and completed less than one circuit of cage before the experimenter caught it.
- Score 2 = The mouse escaped from the first try of the experimenter to catch it and completed 1–2 circuits of cage before the experimenter caught it.
- Score 3 = The mouse escaped from the first try of the experimenter to catch it and completed more than 2 circuits of cage before the experimenter caught it. It happened that the mouse jumped onto the cage wall or the experimenter captured the mouse by the tail on the wall.
- Score 4 = The mouse jumped out of the cage.

## **Social interaction**

After mice were kept in the housing systems for 7 and 11 weeks, the social interaction test was carried out, which was performed in a 43.5 × 43.5 cm arena equipped with a perforated polycarbonate box (10 × 6.5 cm). The approach-avoidance behavior of a test mouse to a C57BL/6JRj mouse was recorded with a video tracking system. The social interaction test consisted of two 2.5-min sessions. In the first 2.5-min 'no target' session, the mouse was allowed to explore the open arena freely with an empty perforated polycarbonate cage. For the second 2.5-min 'target' session, an unfamiliar C57BL/6JRj male mouse was placed into the perforated polycarbonate box. The wire box allowed visual, olfactory, and auditory interactions between the test mouse and the target mouse but prevented direct physical contact. A video tracking software (Ethovision, Noldus, Netherlands) was used to measure the distance moved in the arena and the time spent by the experimental mouse in the "interaction zone" located around the target box (26.0 × 14.5 cm) of the arena. Social interaction was defined as time spent in the interaction zone when the target mouse was present.

## **Anxiety-related behavior**

Trait anxiety-related behavior was examined in the free exploratory paradigm [23,29]. The test was carried out in the testing cage after the burrow had been removed. A gridded cage lid (type I long, 34,5 cm × 14,5 cm) was placed in the testing cage and attached to the back wall using wire. The lid served as a novel object and ladder. Video cameras were installed in a distance of approximately 1.5 m. The mice were video monitored for 5 min and the latency to explore and the total duration of exploration were manually analyzed using ethological analyses software (Etholog version 2.2.5; Ottoni 1999). Experimenters were present during the test and stood silently next to the cameras.

### **Analysis of fecal corticosterone and testosterone metabolites**

Fecal pellets excreted during the period, in which the mice had been individually kept in the testing cage (Makrolon type III cages, 42 × 26 × 15 cm) for approximately three hours (~8.30–11.30 A.M.) to perform the burrowing test and the free exploratory paradigm, were used to analyze fecal corticosterone (FCMs) and testosterone metabolites (FTMs). After this period, testing cages were stored at room temperature for 20–24 hours before fecal pellets were collected from the cages using forceps. FCMs [30-32] and FTMs [33] were extracted and analyzed as previously described.

### **Analysis of hair corticosterone and testosterone**

An electric shaver for small animals (Aesculap Isis GT 420, Suhl, Germany) was utilized to cut off hair samples (approximately 7.5 mg of hair). The first hair sample was taken from the back. Hair took longer than two months to fully regrow, hence there was not enough material for the second sample at the back in week eight. Therefore, the second hair sample was collected from the right hind leg. The third hair sample could be taken from the back, the same body location as the first sample in week twelve of the experiment.

Hair corticosterone and testosterone [pg/mg] were analyzed by liquid chromatography-mass spectrometry in the laboratory of Prof. Kirschbaum, Department of Psychology, Technische Universität Dresden, Germany, as described previously [34].

### **Statistical Analysis**

Statistical analysis was performed with IBM SPSS Version 25 (IBM Corporation, Armonk, NY, USA) and explorative data analysis and tests for normality were performed for each parameter. Differences between the groups were analyzed using the respective test indicated in results section (repeated measures ANOVA, two-way or one-way ANOVA,


Friedman's two-way analysis of variance by ranks, Kruskal-Wallis-Test, Mann-Whitney-U-Test or unpaired Student t-test). Differences were considered significant at  $p < 0.05$ .

## Results

### Burrowing

Since the mice were tested under slightly varying conditions (cage position in the testing room) in the burrowing paradigm, we checked whether the cage position affected the parameters of these tests. Kruskal-Wallis-Test revealed that the cage position had no significant impact on the burrowing performance ( $\text{Chi}^2 = 5.902$ ,  $\text{df} = 5$ ,  $p = 0.316$ ).

Neither on day 1 (Kruskal-Wallis-Test:  $\text{Chi}^2 = 5.295$ ,  $\text{df} = 2$ ,  $p = 0.071$ ), in week 8 ( $\text{Chi}^2 = 2.806$ ,  $\text{df} = 2$ ,  $p = 0.246$ ) nor in week 12 ( $\text{Chi}^2 = 0.433$ ,  $\text{df} = 2$ ,  $p = 0.805$ ) after transfer to the housing systems the burrowing performance differed between housing systems (Figure 3). Friedman's two-way analysis of variance by ranks indicated that burrowing performance increased over time in single-housed mice. In week 12, the re-socialized single/pair mice removed more pellets from the burrow than they did on day 1 in single housing ( $\text{Chi}^2 = 11.375$ ,  $\text{df} = 2$ ,  $z = -3.359$ ,  $p = 0.002$ ). This effect was only found in single/pair-housed mice.




**Figure 3. Burrowing performance.** Data are presented as boxplot diagrams: the box represents the interquartile range (IQR), box edges are the 25th and 75th quartile. The whiskers represent values which are not greater than  $1.5 \times$  IQR. Dots are outliers with values between  $1.5\text{--}3.0 \times$  IQR. Asterisks are outliers with values greater than  $3.0 \times$  IQR. Single housing:  $n = 16$ , group housing:  $n = 12$ , pair housing:  $n = 15$ . Overall, five mice were excluded from statistics because four of them removed less than 5 g food pellets from the burrow (non-responders) and one (group-housed) mouse had to be removed from the experiment due to fight-associated wounds. ##  $p < 0.01$  versus day 1 (Friedman's two-way analysis of variance by ranks).

### Nesting

Nest complexity was only compared between single- and pair-housed mice because nests of group-housed mice had a different shape (nest scores of group-housed mice given as IQR,

median; day 1: 2–4.13, 3.25; week 8: 1.75–4.00, 3.00; week 12: 2.5–3.38, 3.25). On day 1 after transfer to the housing systems, nest scores did not significantly differ between housing systems (Mann-Whitney-U = 92.500,  $z = -1.345$ ,  $p = 0.184$ ; Figure 4A). Eight weeks after mice had been transferred to the housing systems, nests of a higher complexity were found in the pair housing system (Mann-Whitney-U = 189.500,  $z = 2.333$ ,  $p = 0.019$ ). This difference was abolished, when single-housed mice were transferred to pair-housing for 4 weeks (Mann-Whitney-U = 142.000,  $z = 0.539$ ,  $p = 0.616$ ). Friedman's two-way analysis of variance by ranks revealed that nest complexity scores significantly changed over time: both re-socialized single/pair- (versus week 8, in single housing:  $\text{Chi}^2 = 6.933$ ,  $df = 2$ ,  $z = -2.475$ ,  $p = 0.040$ ) and pair-housed (versus day 1:  $\text{Chi}^2 = 18.561$ ,  $df = 2$ ,  $z = -4.066$ ,  $p < 0.001$ ) mice built more complex nests in week 12.


The nest positions of pair-housed mice are illustrated in Figure 4B. Data were analyzed descriptively since mice housed as pairs had to be considered as a unit ( $n = 8$ ) so that the number of units was too low for statistical analysis. Overall, the majority of mice built nests in the units of the cage back under the food unit near the dividing wall. In week 8, two pairs build nests in units bordering one another. There were four nest pairs with a distance of  $\leq 1$  unit from each other and two nest pairs with a distance of 1.5–2 units. Four weeks later, four pairs continuously kept in the pair housing system built nests in units bordering one another, two pairs in units with a distance of  $\leq 1$  unit, and two pairs in units with a distance of 1.5–2 units. Re-socialized single/pair-housed mice showed a slightly different preference for nest location. Only one pair of nest was located next to each other. Three pairs built nests in units with a distance of  $\leq 1$  unit and four pairs with a distance of 1.5–2 units from each other.



**Figure 4: Nesting.** Nest complexity scores (A): Data are presented as boxplot diagrams: the box represents the interquartile range (IQR), box edges are the 25th and 75th quartile. The whiskers represent values which are not greater than  $1.5 \times$  IQR. Dots are outliers with values between  $1.5\text{--}3.0 \times$  IQR. \*  $p < 0.05$  versus pair housing (Kruskal-Wallis-Test), #  $p < 0.01$ , ###  $p < 0.001$  (Friedman's two-way analysis of variance by ranks). Single housing:  $n = 16$ , pair housing:  $n = 16$ . Nest positions in the pair housing system (B): Both the left and right cage compartment were divided into six units. The unit (or border) where the nest was built was determined. Symbols represent a nest; the same symbols were used for nests of both mice of a pair ( $n = 8$  pairs).

### Ease of Handling

Ease of handling was analyzed in week 8. While the latency to first voluntary interaction with the experimenter's hand was significantly higher in pair-housed mice when compared to single-housed mice (Kruskal-Wallis-Test:  $\text{Chi}^2 = 8.483$ ,  $\text{df} = 2$ ,  $z = -2.880$ ,  $p = 0.012$ ; Figure 5), the interaction scores did not differ between single-, group, and pair-housed animals (Kruskal-Wallis-Test:  $\text{Chi}^2 = 2.673$ ,  $\text{df} = 2$ ,  $p = 0.263$ ). The capture score was significantly higher in single-housed mice than in group-housed mice (Kruskal-Wallis-Test:  $\text{Chi}^2 = 14.340$ ,  $\text{df} = 2$ ,  $z = 3.754$ ,  $p = 0.001$ ; Table 1).



**Figure 5: Ease of handling.** Data are presented as boxplot diagrams: the box represents the interquartile range (IQR), box edges are the 25th and 75th quartile. The whiskers represent values which are not greater than  $1.5 \times$  IQR. Single housing:  $n = 16$ , group housing:  $n = 15$ , pair housing:  $n = 16$ . One mouse of the group-housed animals had to be removed from the experiment due to fight-associated wounds. #  $p < 0.05$  versus pair housing (Kruskal-Wallis-Test).

**Table 1. Interaction and capture scores.**

Housing condition	Interaction score	Capture score
Single housing	2 (2–2)	1 (1–0.25)**
Group housing	2 (2–2)	0 (0–0)
Pair housing	2 (2–2)	0.50 (0–1)


Data are given as median (25th quartile – 75th quartile) and analyzed using Kruskal-Wallis-Test: \*\*  $p < 0.01$  versus group housing. Single housing:  $n = 16$ , group housing:



n = 15, pair housing: n = 16. One mouse of the group-housed animals had to be removed from the experiment due to fight-associated wounds.

### **Social interaction**

All mice showed social interaction, reflected by increased time spent in the interaction zone when a target mouse was present ( $F(1, 90) = 9.99$ ,  $p = 0.002$ ; Figure 6), however, social behavior was not affected by the housing condition (repeated measures ANOVA:  $F(2, 90) = 2.57$ ,  $p = 0.82$ ). On the contrary, locomotor activity in the absence of a target mouse differed between housing systems ( $F = (2, 44) = 17.21$ ,  $p < 0.001$ ). The distance moved in the test arena was reduced in pair-housed mice compared to single- (One-way ANOVA:  $F(2, 46) = 10.79$ , Bonferroni post-hoc analysis:  $p < 0.001$ ) and group-housed mice ( $p = 0.025$ ) in week 8. In week 12, both re-socialized mice (single/pair) (One-way ANOVA:  $F(2, 46) = 23.680$ , Bonferroni post-hoc analysis:  $p < 0.001$ ) and pair-housed mice ( $p < 0.001$ ) moved less than group-housed mice.




**Figure 6: Social interaction test.** (A) Time spent in interaction zone in week 8 (left) and week 12 (right). Data are presented as boxplot diagrams: the box represents the interquartile range (IQR), box edges are the 25th and 75th quartile. The whiskers represent values which are not greater than  $1.5 \times \text{IQR}$ . Dots are outliers with values between  $1.5\text{--}3.0 \times \text{IQR}$ . \*  $p < 0.05$ , \*\*\*  $p < 0.001$  compared to group housing; ###  $p < 0.001$  compared to single-housing (one-way ANOVA using Bonferroni post-hoc analysis). Single housing:  $n = 16$ , group housing:  $n = 15$ , pair housing:  $n = 16$ . One mouse of the group-housed animals had to be removed from the experiment due to fight-associated wounds.

### Anxiety-related behavior

Since the mice were tested under slightly varying conditions (cage position in the testing room) in the free exploratory paradigm, we checked whether the cage position affected the parameters of these tests. One-way ANOVA showed that the cage position in the testing room had no significant impact on the logarithm of the latency to explore ( $F(5, 86) = 1.087$ ,  $p$

= 0.373), the logarithm of number of explorations ( $F(5, 86) = 1.064, p = 0.386$ ) or the total duration of exploration ( $F(5, 86) = 1.317, p = 0.265$ ).

Repeated measures ANOVA revealed that there was no effect of time or housing system on the latency to explore (logarithmised) (time:  $F(1, 43) = 3.315, p = 0.076$ ; housing system:  $F(2, 43) = 0.604, p = 0.551$ ) and the duration of exploration within 5 min (time:  $F(1, 43) = 3.055, p = 0.088$ ; housing system:  $F(1, 43) = 1.838, p = 0.171$ ; Figure 7).




**Figure 7: Anxiety-related behavior.** Data are presented as boxplot diagrams: the box represents the interquartile range (IQR), box edges are the 25th and 75th quartile. The whiskers represent values which are not greater than  $1.5 \times$  IQR. Dots are outliers with values between  $1.5-3.0 \times$  IQR. Single housing:  $n = 15$ , group housing:  $n = 15$ , pair housing:  $n = 16$ . One mouse of the group-housed animals had to be removed from the experiment due to fight-associated wounds; one single-housed mouse was excluded because the ladder was not properly attached to the cage wall and slipped down to the cage floor.

### Body weight

During the course of the experiment, a repeated measures ANOVA with a Greenhouse-Geisser correction indicated that there was a significant effect of time on the body weight ( $F(1.95, 85.58) = 133.51, p < 0.001$ ) and a significant interaction between time and housing system ( $F(3.89, 85.58) = 3.25, p = 0.017$ ; Figure 8). Tests of between-subject-effects

revealed that there was a significant difference in body weight between the housing systems ( $F(2, 44) = 4.29, p = 0.020$ ). Dunnett-T3 post-hoc analysis indicated that body weight was significantly higher in group-housed mice when compared to single-housed mice ( $p = 0.02$ ).

The analysis of body weight change in week 9–12 also revealed a significant time effect on the body weight (repeated measures ANOVA with a Greenhouse-Geisser correction:  $F(1.63, 71.92) = 9.61, p < 0.001$ ). Tests of between-subject-effects revealed that there was a significant difference in body weight between the housing systems ( $F(2, 44) = 11.98, p < 0.001$ ). Bonferroni post-hoc analysis indicated that body weight was significantly higher in group-housed mice when compared to re-socialized single (single/pair) mice ( $p < 0.001$ ) and pair-housed mice ( $p = 0.01$ ).



**Figure 8. Body weight.** Data are given as mean  $\pm$  standard deviation. Single housing:  $n = 16$ , group housing:  $n = 15$ , pair housing:  $n = 16$ . One mouse of the group-housed animals had to be removed from the experiment due to fight-associated wounds.

### Analysis of fecal corticosterone and testosterone metabolites

A repeated measure ANOVA was conducted that examined the effect of time and housing system on FCMs or FTMs (Table 2). There was no effect of time (tests of within-subject-effects:  $F(2, 80) = 2.178, p = 0.12$ ) and housing system (tests of between-subject-effects:  $F$

(1, 40) = 0.518,  $p = 0.60$ ) on FCMs, whereas both time (tests of within-subject-effects:  $F(2, 80) = 11.655$ ,  $p < 0.001$ ) and housing system (tests of between-subject-effects  $F(1, 40) = 3.389$ ,  $p = 0.044$ ) significantly affected FTM levels. The post-hoc Bonferroni test revealed that FTMs concentrations were lower on day 1 when compared to week 8 ( $p = 0.003$ ) and week 12 ( $p < 0.001$ ). We analyzed differences in FTM concentrations between housing systems for every time point separately using one-way ANOVA with the post-hoc Bonferroni test: While FTM levels did not significantly differ between housing systems in week 8 ( $F(2, 40) = 0.142$ ,  $p = 0.868$ ) and week 12 ( $F(2, 40) = 2.221$ ,  $p = 0.122$ ), FTM concentrations were significantly lower in group-housed mice when compared to single-housed mice ( $F(2, 40) = 4.582$ ,  $p = 0.016$ ; Bonferroni post-hoc analysis:  $p = 0.048$ ) and pair-housed mice (Bonferroni post-hoc analysis:  $p = 0.023$ ) on day 1.

**Table 2. Analysis of fecal corticosterone and testosterone metabolites concentrations (ng/0.05g)**

Housing system	Day 1	Week 8	Week 12	
<b>Fecal corticosterone metabolites</b>				
Single housing	45.18 ± 18.85	38.91 ± 12.70	Single/Pair housing	30.75 ± 12.84
Group housing	36.30 ± 10.05	42.49 ± 15.29	Group housing	33.29 ± 13.56
Pair housing	36.58 ± 14.87	30.98 ± 12.89	Pair housing	37.06 ± 13.98

Fecal testosterone metabolites in ng/0.05g				
Single housing	3.14 ± 1.04*	3.71 ± 0.70	Single/Pair housing	3.93 ± 1.10
Group housing	2.15 ± 1.16	3.73 ± 0.59	Group housing	3.25 ± 1.22
Pair housing	3.24 ± 0.88*	3.57 ± 1.18	Pair housing	4.22 ± 1.33
<p>Single/pair housing: After week 8, single-housed mice were re-socialized and transferred to the pair-housing system.</p> <p>Data are given as mean ± standard deviation. Single housing: n = 12, group housing: n = 15, pair housing: n = 14. One mouse of the group-housed animals had to be removed from the experiment due to fight-associated wounds; four single-housed mice and two pair-housed mice were excluded because we could not collect enough sample material for analysis within the testing period.</p> <p>* p &lt; 0.05 versus group housing (one-way ANOVA).</p>				

### Analysis of hair corticosterone and testosterone

Hair samples were collected from different body regions (baseline: back, week 8: right leg, week 12: back) due to insufficient hair growth at the back. Therefore, results obtained in week 8 are not comparable with baseline and week 12 and data were analyzed for every time point separately using one-way ANOVA with the post-hoc Bonferroni test (Table 3). Hair corticosterone concentrations did not differ between the study groups at baseline when all mice were still kept in groups (one-way ANOVA:  $F(2, 42) = 0.498$ ,  $p = 0.612$ ). In week 8

( $34.95 \pm 5.89$ ;  $F(2, 42) = 0.616$ ,  $p = 0.545$ ) and week 12 ( $F(2, 42) = 1.230$ ,  $p = 0.303$ ), there were no differences in hair corticosterone concentrations between single-, group-, and pair-housed mice. Repeated measures ANOVA revealed that hair corticosterone concentrations differed between baseline and week 12 ( $F(1, 42) = 232.97$ ,  $p < 0.001$ ) with higher values in week 12 (post-hoc Bonferroni analysis:  $p < 0.001$ ).

Hair testosterone concentrations significantly differed between the study groups at baseline (Welch's  $F(2, 24.606) = 5.278$ ,  $p = 0.012$ ). Therefore, we calculated the percentage change between week 12 and baseline, which was significantly higher in group-housed mice in comparison to pair-housed mice (Kruskal-Wallis-Test:  $\text{Chi}^2 = 15.637$ ,  $\text{df} = 2$ ;  $z = 2.670$ ,  $p = 0.023$ ) and re-socialized mice that were transferred from single to pair housing after week 8 ( $z = -3.860$ ,  $p < 0.001$ ). In week 8, no differences between the housing systems in hair testosterone concentrations were found ( $F(2, 42) = 1.640$ ,  $p = 0.206$ ).

**Table 3. Analysis of hair corticosterone and testosterone concentrations**

Housing system	Baseline (sample location: back; pg/mg)	Week 8 (sample location: right leg; pg/mg)	Week 12 (sample location: back; pg/mg)	%-change between week 12 and baseline
<b>Hair corticosterone</b>				
Single housing	$23.82 \pm 3.77$	$37.72 \pm 6.07$	Single/Pair housing	$34.16 \pm 5.66$
Group housing	$23.84 \pm 4.05$	$36.12 \pm 8.41$	Group housing	$29.30 \pm 8.79$

Pair housing	22.63 ± 3.30	34.95 ± 5.89	Pair housing	32.88 ± 11.59	
<b>Hair testosterone</b>					
Single housing	0.66 ± 0.07*	1.28 ± 0.16	Single/Pair housing	2.07 ± 0.65**	314.47 ± 107.17###
Group housing	0.57 ± 0.09	1.41 ± 0.28	Group housing	2.96 ± 0.99	538.31 ± 224.81
Pair housing	0.66 ± 0.17	1.50 ± 0.50	Pair housing	2.26 ± 0.63	351.01 ± 92.12#

Data are given as mean ± standard deviation. Single housing: n = 16, group housing: n = 15, pair housing: n = 14. One mouse of the group-housed animals had to be removed from the experiment due to fight-associated wounds; two pair-housed mice were excluded because we could not collect enough sample material for analysis within the testing period.

\* p < 0.05 (one-way ANOVA using post-hoc Dunnett T3), \*\* p < 0.01 versus group housing (one-way ANOVA using post-hoc Bonferroni).

# p < 0.05, ### p < 0.001 versus group housing (Kruskal-Wallis-Test).

## Discussion

Since the well-being of laboratory animals significantly contributes to the quality of research [35], it is crucial to understand the impact of housing conditions on the well-being of mice and to further improve them. Both standard housing conditions for male mice, group and single housing, can impair their well-being. Therefore, we are drawing attention to an alternative housing system as a possible refinement strategy: separated pair housing. A perforated



transparent wall divides the cage into two compartments and allows olfactory, acoustical, and visual communication between the two mice but prevents fighting and injuries. In order to enable routine use of separated pair housing in laboratory animal facilities, IVC systems reflecting current international standard housing conditions were used. We systematically assessed the impact of the housing systems (group, single, pair) on the well-being of adult male C57BL/6JRj mice by investigating burrowing and nesting, trait anxiety-related behavior, the ease of handling, social behavior, body weight, and stress hormone (metabolites) concentrations in feces and fur.

The main findings of our study were that the housing system can affect behavioral, biochemical, and physical parameters. Single and pair housing caused initial mild acute stress due to acclimatization to the new housing conditions, though the FTM concentrations must be interpreted with some caution. In week 8, pair-housed mice built more complex nests but showed less locomotor activity and elevated anxiety-related behavior in anticipation of handling. Moreover, differences in body weight were found between single, group, and pair housing with an increased body weight gain in group-housed mice. After single-housed mice were re-socialized and transferred to pair housing, findings in nesting and burrowing suggested that well-being of single-housed mice was improved by transferring them to pair housing.

### **Transfer from group- to pair housing: Long-term effects on well-being**

When mice were 7 weeks old, they were assigned the study groups and hence were transferred from group housing to the respective housing systems (pair, single, group) to be investigated for 8 weeks.

Since nesting and burrowing can be decreased, if well-being of a mouse is reduced [36], we evaluated the long-term effects of the housing systems on both behaviors. Nests of groups had a different structure, i.e. they were wider and flatter, than nests of individuals because up to four mice shared one nest. That is why nest complexity scores were not comparable

between group housing and single or pair housing, but only between single and pair housing. While on day 1 after mice had been transferred to their new housing system, nest complexity did not differ between single- and pair-housed mice, in week 8 pair-housed mice built more complex nests. In contrast, Rettich et al. found nests of a poorer quality when 8–9-month-old vasectomized Hsd:NMRI mice were kept in pairs, separated by a grid [20]. This discrepancy may be explained by differences in age, as previously shown in 25-month-old male C57BL/6 mice that built nests of a lower complexity than 7-month-old animals [37]. Moreover, it has to be noted that our experimental design deviated from Rettich et al.'s, which could have a potential effect on the well-being of the mice. First, the habituation period to the housing system was much shorter in Rettich et al. (9 days) [20] than in our study (8 weeks). Secondly, in contrast to our study, Rettich et al.'s mice had already experienced previous experimental phases (i.e. 3 × 14-day period of single-housing followed by rehousing with a female for several weeks) before they were transferred to the pair housing system with an unfamiliar male. This may have enhanced territorial authority as well as aggressive behavior [20] causing higher distress levels than in mice of our study. Thirdly, differences in cage ventilation must be considered because the movement of air causes heat loss through convection and, therefore, can affect nest building behavior [38,39]. High nest walls can protect mice from the draught caused by high ventilation rates [39]. For instance, female C57BL/6NCrl mice built more complex nests when they were kept in IVCs with 75 air changes per hour when compared to IVCs with 50 air changes per hour [38]. With regard to our study, the ventilation rate differed between IVCs used pair housing (75 changes per hour) and IVCs used for single housing (50–60 changes per hour), which could explain the higher nest complexity in pair-housed mice. Another important issue is that the experimenter assessing the nests could not be blinded because the pair housing system with its cage divider looks different than the cages used for single housing, even on images or videos.

To investigate whether mice could meet their need for proximity in the pair housing system, we determined the positions of the nest sites. Mice obviously preferred to build nests in the rear third of the cage under the food unit, probably due to a lower light intensity of 16 lx in

the average in this part of the cage compared to 40 lx on the front side. Moreover, in the back of the cage under the air inlet the draught may be less, as dead-air spaces are created [39]. We found that the majority of pairs located their nests with a distance of  $\leq 1$  unit from each other and, interestingly, most nests were positioned next to the cage divider, which made us to hypothesize that the mice preferred the proximity to the other male rather than staying alone. However, this may also be due to the intention of sharing body heat. It was shown earlier that male mice choose dwelling next to a familiar cage mate [19], for instance nests of pair-housed male mice were located close to each other [20].

Mice are highly social animals and usually interact with unfamiliar mice intruding into their territories. Social interaction of mice appeared not to be affected by the housing systems, as demonstrated in the social interaction test using a male mouse as novel intruder, though results are difficult to interpret because the distance moved in the arena differed between housing conditions. Interestingly, mice kept as pairs moved less than single- and group-housed animals which may indicate a decrease in exploratory behavior and is why the time spent in the interaction zone should be interpreted with caution. A higher locomotor activity in socially isolated C57BL/6 mice when compared to group housing was previously found in behavioral paradigms such as the open field [16,17]; pair housing seemed to have the opposite effect. Locomotor activity can be affected by stress [40-44], hence it may indicate higher stress levels in single-housed mice. We did not investigate whether these effects are also present in the home cage, i.e. home cage activity and time spent near the dividing wall should be further investigated. Moreover, it would be of high interest to monitor whether and how pair-housed mice actually interact with the other mouse beyond the dividing wall and whether the dividing wall is marked with urine to indicate territorial boundaries more intensive than other areas of the cage.

Considering the impact of the housing systems on locomotor activity, results obtained from the free exploratory paradigm also need to be carefully interpreted. Locomotor activity could not be determined in this test. In the free exploratory paradigm, all mice, independently of the housing system, explored the gridded cage lid and no effects of the housing systems on trait

anxiety-related behavior was detected. In contrast, a decrease in trait anxiety was found in 12-week old male Swiss CD-1 mice that had been socially isolated for 3 weeks when compared to group-housed animals [14]. Moreover, changes in state anxiety-related behavior induced by social isolation, depending on mouse strain, age, and duration of isolation, were reported. While state anxiety in the elevated plus maze test increased in 7-week old male ICR mice that had been single-housed for 4 weeks [45] in comparison to group-housed mice, it decreased in 11-week-old male DBA and C57BL/6J mice (7 weeks of social isolation) [16].

To investigate whether the housing systems influenced the human-animal interaction and the ease of handling during routine husbandry procedures, we measured the latency to first voluntary interaction with the experimenter's hand and used an interaction as well as a capture scoring system. While all mice, independently of the housing systems, explored the experimenter's hand by direct contact of whiskers and/or nose, pair-housed mice showed a prolonged latency to first voluntary interaction when compared to single-housed mice. This may be explained by their lower locomotor activity we found in the social interaction test. Hurst and West [27] previously used the latency to first voluntary interaction with the experimenter's hand to assess the impact of tail, tunnel, and cup handling on the "anxiety-related behavior in anticipation of handling". Mice used in the present study were tail handled. There was one striking difference between the cages that were used for the different housing systems: for handling the mice, the food unit had to be removed from the cages that were used for single and group housing (i.e. Polysulfone type I and II long), whereas the food unit of the Green Line IVC Sealsafe PLUS Rat cages, our pair housing system, remained in the same position. When pair-housed mice had to be removed from the cages for routine husbandry procedures, they clung tightly to the grid of the food unit, our animal care technicians reported. One can imagine that more strength must be expended when picking up a mouse clinging to a grid than a mouse sitting on the cage floor, which may be why pair-housed mice had worse handling experiences than single- and group-housed animals and showed higher anxiety-related behavior in anticipation of handling. The potential

impact of the food unit providing hold for the mice was an incidental finding and was only revealed due to the good communication between technicians and scientists. Since it was not considered in the experimental design of the study, further investigations are needed to prove our interpretation of these results. The capture scores did not reflect the observations of our technicians, because the food unit was removed for this test to create equal conditions for all experimental groups. Single-housed mice were more difficult to catch than group-housed mice. Single-housed mice may be less used to moving objects in their environment because they do not share their cages with other animals. The moving hand of the experimenter may trigger their flight response to a higher extent than in pair- and group-housed mice.

FCM and hair corticosterone concentrations were used as markers for stress. Since the excretion of FCMs depends on the circadian rhythm [31], fecal samples were always collected at the same time of day (~8.30–11.30 A.M.). The results reflected the hypothalamic-pituitary-adrenal axis activity 8–10 hours before [31], i.e. in the dark period on day 1 or week 8 after transfer to the housing systems. Hair corticosterone may not only indicate ongoing stress [46] the mice experienced at the time when the samples were taken but also serve as a retrospective biomarker for stress, as corticosterone can accumulate in hair over time [47]. The analysis of FCMs and hair corticosterone concentrations did not reveal any significant short- and long-term effects of the housing systems on the stress hormone (metabolites) levels. Hair corticosterone increased over time, as expected [47]. Pair housing appeared not to be less stressful than single or group housing, as living next to a male rival may be as stressful as social isolation or being part of a hierarchy [48]. This is in line with other findings, which neither found changes in FCMs in male C57BL/6J mice [49] nor in plasma corticosterone concentrations in males CD-1 [14], male TO albino mice [50] or male Swiss albino mice [51] when the animals were separated. In contrast, a transient increase in urine corticosterone concentrations was found a day after male C57BL/6NCRI mice were transferred to individual cages, however, their urine corticosterone concentrations were lower in comparison to group-housed mice on day 7 [10]. When analyzing hormone

(metabolites) levels, the social context and social rank of the individuals should also be considered to investigate the correlation between the hormone (metabolite) levels and the social behavior of the mice [52]. Plasma corticosterone levels were demonstrated to be lower in alpha male CD-1 mice than in subordinates in despotic groups [52]. Stress levels of the latter were shown to be higher in larger groups than in groups of two mice only [52]. Against this background, it would be of high interest to further investigate whether mice establish a hierarchy in the pair housing system and whether the lack of a clear hierarchy, which probably is associated with ongoing social defeat through the cage divider, may be a reason for stress levels comparable with single and group housing.

Besides corticosterone (metabolites), testosterone is also dependent on the group structures and the individual social ranks [52]. Testosterone levels of alpha male mice in despotic groups are higher than in subordinates. A flat hierarchy and also a victory result in higher testosterone concentrations of subordinate male mice when compared to despotic group behavior and defeat [52,53]. In the initial stage of acute stress, testosterone levels can be elevated; a variety of factors (e.g. the absence of chronic stress, dominant status) contribute to this transient increase [53]. On day 1, FTM levels were elevated in single- and pair-housed mice when compared to group-housed animals, suggesting mild acute stress due to acclimatization to the new housing condition and, for pair-housed mice, to the unfamiliar cage mate. In the latter case, the increase in testosterone levels may be due to the agonist encounter with the other male mouse. As group-housed mice were siblings, it is likely that the hierarchy in the group was flat and fights for rank were rare. In week 8, the mice seemed to have recovered from this initial stress and habituated the housing conditions; their FTM concentrations were increased when compared to day 1. It is known that chronic stress decreases testosterone (serum) concentrations [54], which in turn indicates that neither of the housing system caused chronic stress.

However, hair testosterone concentrations differed between the three study groups at baseline, so it is possible that testosterone values had already been higher in mice that were assigned to the single and pair housing group before they were transferred to the housing

systems. Therefore, both hair as well as fecal testosterone values should be interpreted with caution. Since hair samples were taken from different body parts (i.e. from the back at baseline, from the right hind leg in week 8), it is not clear to the authors if values can be compared between these two time points. Hair corticosterone concentrations may depend on the sample location.

Body weight was significantly lower in single- and pair-housed mice compared to group-housed mice. A difference in body weight between single and group housing had already been demonstrated in male C57BL/6J [9,16] and male C57BL/6JRj [17]. Along with this, it would be interesting to investigate whether the weight in single- and pair-housed mice decreased due to reduced food intake, higher activity in the home cage or enhanced thermoregulation. With regard to the latter point, single- and pair-housed mice may need more energy to maintain body temperature because they cannot huddle together to keep warm while resting.

Overall, we could not identify clear long-term beneficial effects of pair housing on the well-being of male C57BL6/JRj mice.

### **Transfer from single to pair housing**

At the age of 15 weeks, single-housed mice were transferred to the pair-housing system in order to re-socialize them (hereafter also referred to as single/pair housing). The effects of the transfer on their well-being were investigated after three to four weeks, i.e. in week 11 and 12 of the study.

Body weight remained higher in group-housed mice and no changes in trait anxiety-related behavior, and hormone (metabolite) concentrations, except from hair testosterone, were found. Hair testosterone concentrations increased in all study groups, but the percentage change to baseline was lower in (re-socialized) pair-housed mice in comparison to group-housed animals. These findings suggested that higher levels of testosterone were

incorporated into the hair shaft in group-housed mice over the experimental period, which may be explained by the dominance hierarchy and the fights involved, resulting in high testosterone levels in dominant mice and most attacked subordinate animals [55]. However, concentrations of fecal testosterone metabolites (FTMs) did not reflect this hypothesis. The discrepancies may be explained by the low number of sampling time points, which did not allow to determine the excretion of FTMs over the whole experimental period but only at certain time points. In contrast, the level of testosterone measured at a certain time can reflect several weeks of hormone accumulation [47], which may give more information on the hormonal balance of the mice in this case.

Nest complexity and burrowing performance suggested that the well-being of the re-socialized mice increased under the pair-housing conditions. While nest complexity was poorer in single- than pair-housed mice in week 8, there were no differences in week 12 anymore, as nest complexity significantly increased over time. This may indicate fostered well-being of single/pair-housed mice, but can also be explained by the different ventilation rates with a higher movement causing heat loss through convection, which made the mice to build nests with higher walls [38,39]. In contrast to pair-housed mice, the majority of re-socialized mice appeared not to prefer to a nest location next to the cage divider. Rettich et al. [20] explained the short-term negative effects of male:male pair housing in 8–9-month-old vasectomized Hsd:NMRI mice by stronger territorial authority, which may also influence their choice of nest locations. However, based on our data, we cannot conclude with certainty that the nest locations are a sign for a stronger territorial authority due to the 8-week period of single housing. Further investigations are needed to prove this hypothesis, including the time spent near the dividing wall, the interaction through the cage divider, and the urine marks indicating their territorial boundaries.

Although burrowing performance did not differ between single/pair, group, and pair housing in week 12, re-socialized mice removed more pellets from the burrow when compared to day 1. Whether this is due to a habituation effect to the burrowing test or due to improved well-



being is unclear, since FCM concentrations did not reveal any time-dependent changes in stress levels.

In week 11, mice were tested again in the social interaction test. Interestingly, re-socialization of single-housed mice reduced the locomotor activity the arena when compared to group-housed mice and reaches a level comparable to mice that had been kept in the pair housing systems since day 1 of the study. These findings confirmed the results we found in week 8 and again raised the question whether home cage activity was also affected. This should be tackled in future studies, as stated above, in order to learn more about the underlying reasons. It also would be of high interest to determine the interaction frequency and quality between the re-socialized cage companions along the dividing wall and compare their behavior with mice that were directly transferred from group to pair housing. Moreover, it remains unknown whether anxiety-related behavior in anticipation of handling increased and capture scores decreased in re-socialized mice because, unfortunately, the ease of handling was not tested in week 12 again.

All in all, the re-socialization seemed to improve well-being of single-housed mice to a slight extent. This hypothesis is supported by An et al. demonstrating that male KM and BALB/c mice recovered from single housing induced effects (i.e cognitive effects) [56].

### **Future work**

To clarify whether pair housing is an appropriate refinement measure for male mice, further investigations are needed, including the interaction through the cage divider, urine marks for delineating territorial boundaries (i.e. the divider), and the time spent close to each other. Moreover, the effects of the housing systems on the affective state of the mice should be investigated more profoundly using tests for state anxiety-related behavior as well as a cognitive bias test.

## **Conclusion**

The results of the present study did not reveal unambiguous long-term beneficial effects of separated pair housing on the well-being of adult male C57BL/6JRj mice, though the transfer from single to pair housing appeared to slightly foster well-being with regard to nest complexity and burrowing. Taking into account that male mice prefer dwelling near other males to staying alone, pair housing rather than single housing can meet this need, as reflected by the nest locations. However, neither single nor group housing of adult male C57BL/6JRj mice appeared to have a severe impact on the well-being of the mice. It should be noted that the effects of the housing systems on behavioral, physical, and biochemical parameters needs to be considered in the design of animal experimental studies and their analysis.

## References

- 1 Olsson, I. A. S. & Westlund, K. More than numbers matter: The effect of social factors on behaviour and welfare of laboratory rodents and non-human primates. *Appl Anim Behav Sci.* **103**, 229-254 (2007).
- 2 National Research Council (US). Guide for the Care and Use of Laboratory Animals. . Available from: <https://grants.nih.gov/grants/olaw/guide-for-the-care-and-use-of-laboratory-animals.pdf> (last accessed on 1 March 2019). (2001).
- 3 Council of Europe. Appendix A of the European Convention for the Protection of Vertebrate Animals Used for Experimental and Other Scientific Purposes (ETS No. 123). In: Guidelines for Accommodation and Care of Animals. Strasbourg, France: CE (2006).
- 4 Directive 2010/63/EU of the European Parliament and of the council of 22 September 2010 on the protection of animals used for scientific purposes. *Official Journal of the European Union ABI.* **L276/33-L276/79** (2010).
- 5 Kappel, S., Hawkins, P. & Mendl, M. To group or not to group? Good practice for housing male laboratory mice. *Animals (Basel).* **7**, 88 (2017).
- 6 Weber, E. M., Dallaire, J. A., Gaskill, B. N., Pritchett-Corning, K. R. & Garner, J. P. Aggression in group-housed laboratory mice: why can't we solve the problem? *Lab Animal (NY).* **46**, 157-161 (2017).
- 7 Van Loo, P. L. P., Van Zutphen, L. & Baumans, V. Male management: coping with aggression problems in male laboratory mice. *Lab Anim.* **37**, 300-313 (2003).
- 8 Tallent, B. R., Law, L. M., Rowe, R. K. & Lifshitz, J. Partial cage division significantly reduces aggressive behavior in male laboratory mice. *Lab Anim.* **52**, 384-393 (2018).
- 9 Nagy, T. R., Krzywanski, D., Li, J., Meleth, S. & Desmond, R. Effect of group vs. single housing on phenotypic variance in C57BL/6J mice. *Obes Res.* **10**, 412-415 (2002).
- 10 Kamakura, R., Kovalainen, M., Leppäluoto, J., Herzig, K. H. & Mäkelä, K. A. The effects of group and single housing and automated animal monitoring on urinary corticosterone levels in male C57BL/6 mice. *Physiol Rep.* **4**, e12703, doi:10.14814/phy2.12703 (2016).
- 11 Chourbaji, S., Zacher, C., Sanchis-Segura, C., Spanagel, R. & Gass, P. Social and structural housing conditions influence the development of a depressive-like phenotype in the learned helplessness paradigm in male mice. *Behav Brain Res.* **164**, 100-106 (2005).
- 12 Kallioikoski, O., Teilmann, A. C., Jacobsen, K. R., Abelson, K. S. & Hau, J. The lonely mouse—single housing affects serotonergic signaling integrity measured by 8-OH-DPAT-induced hypothermia in male mice. *PLoS One.* **9**, e111065 (2014).
- 13 Macedo, G. C. *et al.* Consequences of continuous social defeat stress on anxiety-and depressive-like behaviors and ethanol reward in mice. *Horm Behav.* **97**, 154-161 (2018).
- 14 Bartolomucci, A. *et al.* Individual housing induces altered immuno-endocrine responses to psychological stress in male mice. *Psychoneuroendocrinology.* **28**, 540-558 (2003).
- 15 Berry, A. *et al.* Social deprivation stress is a triggering factor for the emergence of anxiety-and depression-like behaviours and leads to reduced brain BDNF levels in C57BL/6J mice. *Psychoneuroendocrinology.* **37**, 762-772 (2012).
- 16 Voikar, V., Polus, A., Vasar, E., Rauvala, H. Long-term individual housing in C57BL/6J and DBA/2 mice: assessment of behavioral consequences. *Genes Brain Behav.* **4**, 240-252 (2005).
- 17 Pasquarelli, N., Voehringer, P., Henke, J. & Ferger, B. Effect of a change in housing conditions on body weight, behavior and brain neurotransmitters in male C57BL/6J mice. *Behav Brain Res.* **333**, 35-42 (2017).
- 18 Van Loo, P. L., de Groot, A. C., Van Zutphen, B. F. & Baumans, V. Do male mice prefer or avoid each other's company? Influence of hierarchy, kinship, and familiarity. *J Appl Anim Welf Sci.* **4**, 91-103 (2001).
- 19 Van Loo, P. L., Van de Weerd, H. A., Van Zutphen, L. & Baumans, V. Preference for social contact versus environmental enrichment in male laboratory mice. *Lab Anim.* **38**, 178-188 (2004).

- 20 Rettich, A., Käsermann, H. P., Pelczar, P., Bürki, K. & Arras, M. The physiological and behavioral impact of sensory contact among unfamiliar adult mice in the laboratory. *J Appl Anim Welf Sci.* **9**, 277-288 (2006).
- 21 Deacon, R. M. Burrowing in rodents: a sensitive method for detecting behavioral dysfunction. *Nat Protoc.* **1**, 118-121 (2006).
- 22 Deacon, R. M., Raley, J. M., Perry, V. H. & Rawlins, J. N. Burrowing into prion disease. *Neuroreport* **12**, 2053-2057 (2001).
- 23 Hohlbaum, K. *et al.* Systematic assessment of well-being in mice for procedures using general anesthesia. *J Vis Exp* 2018. 133, e57046; 10.3791/57046 (2018).
- 24 Jirkof, P. *et al.* Burrowing behavior as an indicator of post-laparotomy pain in mice. *Front Behav Neurosci.* **4**, 165, 10.3389/fnbeh.2010.00165 (2010).
- 25 Gaskill, B. N., Karas, A. Z., Garner, J. P. & Pritchett-Corning, K. R. Nest building as an indicator of health and welfare in laboratory mice. *J Vis Exp.* 82, 51012; 10.3791/51012 (2013).
- 26 Hess, S. E. *et al.* Home improvement: C57BL/6J mice given more naturalistic nesting materials build better nests. *J Am Assoc Lab Anim Sci.* **47**, 25-31 (2008).
- 27 Hurst, J. L. & West, R. S. Taming anxiety in laboratory mice. *Nat Meth.* **7**, 825-826, doi:10.1038/nmeth.1500 (2010).
- 28 Wahlsten, D., Metten, P. & Crabbe, J. C. A rating scale for wildness and ease of handling laboratory mice: results for 21 inbred strains tested in two laboratories. *Genes Brain Behav.* **2**, 71-79 (2003).
- 29 Bert, B., Schmidt, N., Voigt, J. P., Fink, H. & Rex, A. Evaluation of cage leaving behaviour in rats as a free choice paradigm. *J Pharmacol Toxicol Methods.* **68**, 240-249, 10.1016/j.vascn.2013.01.001 (2013).
- 30 Palme, R., Touma, C., Arias, N., Dominchin, M. F. & Lepschy, M. Steroid extraction: get the best out of faecal samples. *Wien Tierarz Monats* **100**, 238-246 (2013).
- 31 Touma, C., Palme, R. & Sachser, N. Analyzing corticosterone metabolites in fecal samples of mice: a noninvasive technique to monitor stress hormones. *Horm Behav* **45**, 10-22 (2004).
- 32 Touma, C., Sachser, N., Mostl, E. & Palme, R. Effects of sex and time of day on metabolism and excretion of corticosterone in urine and feces of mice. *Gen Comp Endocrinol.* **130**, 267-278 (2003).
- 33 Auer, K. E. *et al.* Measurement of fecal testosterone metabolites in mice: replacement of invasive techniques. *Animals.* **10**, 10.3390/ani10010165 (2020).
- 34 Gao, W. *et al.* Quantitative analysis of steroid hormones in human hair using a column-switching LC-APCI-MS/MS assay. *J Chromatogr B Analyt Technol Biomed Life Sci.* **928**, 1-8, 10.1016/j.jchromb.2013.03.008 (2013).
- 35 Poole, T. Happy animals make good science. *Lab Anim.* **31**, 116-124 (1997).
- 36 Jirkof, P. Burrowing and nest building behavior as indicators of well-being in mice. *J Neurosci Methods.* **234**, 139-146, 10.1016/j.jneumeth.2014.02.001 (2014).
- 37 Xiong, X.-D., Xiong, W.-D., Xiong, S.-S. & Chen, G.-H. Age- and gender-based differences in nest-building behavior and learning and memory performance measured using a radial six-armed water maze in C57BL/6 Mice. *Behav Neurol.* **2018**, 8728415: 10.1155/2018/8728415 (2018).
- 38 Spangenberg, E., Wallenbeck, A., Eklöf, A., Carlstedt-Duke, J. & Tjäder, S. Housing breeding mice in three different IVC systems: maternal performance and pup development. *Lab Anim.* **48**, 193-206 (2014).
- 39 Baumans, V., Schlingmann, F., Vonck, M. & van Lith, H. A. Individually ventilated cages: beneficial for mice and men? *Contemp Top Lab Anim Sci.* **41**, 13-19 (2002).
- 40 Grootendorst, J., De Kloet, E., Vossen, C., Dalm, S. & Oitzl, M. J. Repeated exposure to rats has persistent genotype-dependent effects on learning and locomotor activity of apolipoprotein E knockout and C57Bl/6 mice. *Behav Brain Res.* **125**, 249-259 (2001).
- 41 Strekalova, T., Spanagel, R., Dolgov, O. & Bartsch, D. Stress-induced hyperlocomotion as a confounding factor in anxiety and depression models in mice. *Behav Pharmacol.* **16**, 171-180 (2005).

- 42 Lee, E. H., Tang, Y. P. & Chai, C. Y. Stress and corticotropin-releasing factor potentiate center  
region activity of mice in an open field. *Psychopharmacology (Berl)*. **93**, 320-323 (1987).
- 43 Lee, E. H., Tsai, M. J., Chai, C. Y. Stress selectively influences center region activity of mice in  
an open field. *Physiol Behav*. **37**, 659-662 (1986).
- 44 Zhu, S., Shi, R., Wang, J., Wang, J.-F. & Li, X.-M. Unpredictable chronic mild stress not chronic  
restraint stress induces depressive behaviours in mice. *Neuroreport*. **25**, 1151-1155 (2014).
- 45 Koike, H. *et al.* Behavioral abnormality and pharmacologic response in social isolation-reared  
mice. *Behav Brain Res*. **202**, 114-121 (2009).
- 46 Kalliokoski, O., Jellestad, F. K. & Murison, R. A systematic review of studies utilizing hair  
glucocorticoids as a measure of stress suggests the marker is more appropriate for  
quantifying short-term stressors. *Sci Rep*. **9**, 1-14 (2019).
- 47 Carlitz, E. H. D. *et al.* Steroid hormones in hair reveal sexual maturity and competition in wild  
house mice (*Mus musculus domesticus*). *Sci Rep*. **9**, 16925, 10.1038/s41598-019-53362-4  
(2019).
- 48 Krohn, T. C., Sørensen, D. B., Ottesen, J. L. & Hansen, A. K. The effects of individual housing  
on mice and rats: a review. *Animal Welfare*. **15**, 343-352 (2006).
- 49 Melotti, L. *et al.* Can live with 'em, can live without 'em: Pair housed male C57BL/6J mice  
show low aggression and increasing sociopositive interactions with age, but can adapt to  
single housing if separated. *Appl Anim Behav Sci*. **214**, 79-88 (2019).
- 50 Goldsmith, J. F., Brain, P. F., Benton, D. Effects of the duration of individual or group housing  
on behavioural and adrenocortical reactivity in male mice. *Physiol Behav*. **21**, 757-760  
(1978).
- 51 Misslin, R., Herzog, F., Koch, B. & Ropartz, P. Effects of isolation, handling and novelty on the  
pituitary-adrenal response in the mouse. *Psychoneuroendocrinology*. **7**, 217-221 (1982).
- 52 Williamson, C. M., Lee, W., Romeo, R. D., Curley, J. P. Social context-dependent relationships  
between mouse dominance rank and plasma hormone levels. *Physiol Behav*. **171**, 110-119  
(2017).
- 53 Chichinadze, K., Chichinadze, N. Stress-induced increase of testosterone: contributions of  
social status and sympathetic reactivity. *Physiol Behav*. **94**, 595-603 (2008).
- 54 Son, S. W. *et al.* Testosterone depletion increases the susceptibility of brain tissue to  
oxidative damage in a restraint stress mouse model. *J Neurochem*. **136**, 106-117 (2016).
- 55 Van Loo, P. *et al.* Influence of cage enrichment on aggressive behaviour and physiological  
parameters in male mice. *Appl Anim Behav Sci*. **76**, 65-81 (2002).
- 56 An, D. *et al.* Effects of social isolation, re-socialization and age on cognitive and aggressive  
behaviors of Kunming mice and BALB/c mice. *Anim Sci*. **88**, 798-806 (2017).

## **Acknowledgements**

We thank PD Dr. med. vet. habil. Roswitha Merle and Dr. rer. nat. Sopia K. Piper for assisting with statistical analysis and Larissa Mosch for performing the social interaction test.

## **Author Contributions**

Conceptualization: SF, AR, CTR, KU; Project Administration: KU; Funding Acquisition: KU; Investigation: KH, KU; Data analysis: KH, SF, RP, KU; Visualization: KH, KU; Writing – Original Draft Preparation: KH, KU; Writing – Review & Editing: KH, SF, AR, RP, CTR, KU

## **Competing interests**

The authors declare no competing interests.

## **Funding**

This research was financially supported by the Charité 3R, Charité - Universitätsmedizin Berlin. The funder had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.